



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2006

Is the association of alcohol use disorders with major depressive disorder a consequence of undiagnosed bipolar-II disorder?

Angst, J ; Gamma, A ; Endrass, J ; Rössler, W ; Ajdacic-Gross, V ; Eich, D ; Herrell, R ; Merikangas, K R

Abstract: BACKGROUND: There is emerging evidence that there is a spectrum of expression of bipolar disorder. This paper uses the well-established patterns of comorbidity of mood and alcohol use disorder to test the hypothesis that application of an expanded concept of bipolar-II (BP-II) disorder might largely explain the association of alcohol use disorders (AUD) with major depressive disorder (MDD). **METHOD:** Data from the Zurich study, a community cohort assessed over 6 waves from ages 20/21 to 40/41, were used to investigate the comorbidity between mood disorders and AUD. Systematic diagnostic criteria were used for alcohol abuse, alcohol dependence, MDD, and BP-II. In addition to DSM criteria, two increasingly broad definitions of BP-II were employed. **RESULTS:** There was substantially greater comorbidity for the BP-II compared to major depression and for alcohol dependence compared to alcohol abuse. The broadest concept of BP-II explained two thirds of all cases of comorbidity of AUD with major depressive episodes (MDE). In fact, the broader the definition of BP-II applied, the smaller was the association of AUD with MDD, up to non-significance. In the majority of cases, the onset of bipolar manifestations preceded that of drinking problems by at least 5 years. **CONCLUSIONS:** The findings that the comorbidity of mood disorders with AUD was primarily attributable to BP-II rather than MDD and that bipolar symptoms usually preceded alcohol problems may encourage new approaches to prevention and treatment of AUD.

DOI: <https://doi.org/10.1007/s00406-006-0673-3>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-49810>

Journal Article

Published Version

Originally published at:

Angst, J; Gamma, A; Endrass, J; Rössler, W; Ajdacic-Gross, V; Eich, D; Herrell, R; Merikangas, K R (2006). Is the association of alcohol use disorders with major depressive disorder a consequence of undiagnosed bipolar-II disorder? *European Archives of Psychiatry and Clinical Neuroscience*, 256(7):452-457.

DOI: <https://doi.org/10.1007/s00406-006-0673-3>

ORIGINAL PAPER

Jules Angst · Alex Gamma · Jérôme Endrass · Wulf Rössler · Valdet Ajdacic-Gross · Dominique Eich
Richard Herrell · Kathleen Ries Merikangas

Is the association of alcohol use disorders with major depressive disorder a consequence of undiagnosed bipolar-II disorder?

Received: 2 September 2005 / Accepted: 4 April 2006 / Published online: 17 August 2006

Abstract *Background* There is emerging evidence that there is a spectrum of expression of bipolar disorder. This paper uses the well-established patterns of comorbidity of mood and alcohol use disorder to test the hypothesis that application of an expanded concept of bipolar-II (BP-II) disorder might largely explain the association of alcohol use disorders (AUD) with major depressive disorder (MDD). *Method* Data from the Zurich study, a community cohort assessed over 6 waves from ages 20/21 to 40/41, were used to investigate the comorbidity between mood disorders and AUD. Systematic diagnostic criteria were used for alcohol abuse, alcohol dependence, MDD, and BP-II. In addition to DSM criteria, two increasingly broad definitions of BP-II were employed. *Results* There was substantially greater comorbidity for the BP-II compared to major depression and for alcohol dependence compared to alcohol

abuse. The broadest concept of BP-II explained two thirds of all cases of comorbidity of AUD with major depressive episodes (MDE). In fact, the broader the definition of BP-II applied, the smaller was the association of AUD with MDD, up to non-significance. In the majority of cases, the onset of bipolar manifestations preceded that of drinking problems by at least 5 years. *Conclusions* The findings that the comorbidity of mood disorders with AUD was primarily attributable to BP-II rather than MDD and that bipolar symptoms usually preceded alcohol problems may encourage new approaches to prevention and treatment of AUD.

Key words alcohol use disorders · alcohol abuse · alcohol dependence · bipolar II disorder · major depression · comorbidity

J. Angst (✉) · A. Gamma · W. Rössler
Zurich University Psychiatric Hospital
Lenggstrasse 31
Mail Box 1931
8032 Zurich, Switzerland
Tel.: +41-1/384-2611
Fax: +41-1/384-2446
E-Mail: jangst@bli.unizh.ch

V. Ajdacic-Gross · D. Eich
Zurich University Psychiatric Hospital
Social Psychiatry Research Group
Zurich, Switzerland

R. Herrell · K.M. Merikangas
Section on Developmental Genetic Epidemiology
Intramural Research Program
National Institute of Mental Health
National Institutes of Health
Department of Health and Human Services
Bethesda, MD, USA

J. Endrass
Psychiatrisch-Psychologischer Dienst
Justizvollzug Kanton Zürich
Switzerland

Introduction

There is emerging evidence from prospective studies of community and clinical samples that a substantial proportion of individuals with lifetime major depression may in fact manifest bipolar disorder [1–3, 5, 17, 20, 21]. Application of a broader concept of bipolar disorder (i.e., by including overactivity as one of the core criteria rather than mood elevation or irritability as the sole criterion, and omitting a duration threshold) suggests that approximately 50% of all mood disorders may be more appropriately classified as bipolar II disorder [5]. Rates of major depression may be artificially elevated because of the delayed recognition of hypomanic symptoms as pathological, under-reporting of hypomanic symptoms, which subjects experience as normal, and the high threshold for the diagnosis of hypomania in the DSM-IV [5]. This has relevance for our approach to investigating the association with alcohol use disorders.

Patterns of co-occurrence of alcoholism and mood disorders (both MDD and BP disorders) have been well-established in clinical [10, 30] and community samples [3–5]. The magnitude of the association between mood and alcohol use disorders (AUD) differs substantially according to subtypes of both disorders. For example, alcohol dependence has been found to be far more strongly associated with mood disorders and with most other comorbid conditions, than either problematic alcohol use or abuse [23].

Convergent evidence from large-scale population-based studies has demonstrated a greater association of alcohol dependence [7, 12, 19, 28] with bipolar disorder than with major depressive disorder. Moreover, the NCS study reported that alcohol dependence-mania comorbidity was greater in males than in females (e.g., OR = 12.0 for males, OR = 5.0 for females) [19]. In one of the largest samples to date, Grant et al. [15] confirmed that the association of AUD was consistently higher with mania and hypomania than with MDD or dysthymia [16]. The clinical significance of alcohol use-mood disorder comorbidity is reflected in the increased prevalence of bipolar disorder among alcohol-dependent subjects who had attempted suicide [26, 27].

Therefore, the major aim of this study is to test whether lowering bipolar-II (BP-II) diagnostic thresholds will lead to a reduction in comorbidity between major depression and alcoholism. Further, we propose that a stepwise broadening of the diagnostic criteria for bipolar disorder would not weaken its association with AUD.

Methods

Sample

The Zurich study sample was selected from 4,547 subjects (2,201 males, 2,346 females) of age 19/20, representative of the canton of Zurich in Switzerland. These subjects were screened in 1978 with the Symptom Checklist 90-R (SCL-90-R) [11] in order to oversample persons with high global severity indices (GSI) who would be more likely to develop psychiatric syndromes. A stratified random sample of 591 subjects (292 males, 299 females) was selected for interview and subsequent follow-up. Two-thirds of the cohort consisted of high scorers (defined as those with scores > the 85th percentile on the GSI). The remaining third were a random sample of those with scores below the 85th percentile. The screening took

place in 1978 when the male participants were 19 and the females 20 years of age. Six interview waves were conducted across 20 years, as shown in the Table 1. The interviews assessed a broad range of psychiatric and somatic syndromes and symptoms, including questions about duration, frequency and treatment as well as subjective impairment and distress. Across 20 years the attrition rate was 38% (men 44%, women 31%); 91% of the subjects were interviewed at least twice.

Diagnostic criteria

Diagnoses were derived by computer algorithm from the answers given in the interviews. Alcohol abuse and dependence were classified according to DSM-IV criteria (1994). This retrospective classification was possible because the interview's assessment of alcohol consumption and its consequences was not limited to diagnostic criteria. Major depressive episodes (MDE) were defined by DSM-III criteria (in the first two interviews 1979 and 1981) and DSM-III R criteria (1987) in the last four interviews (1986, 1988, 1993, 1999). Mania was defined according to DSM-IV criteria. BP-II disorder required the presence of a MDE plus hypomania. Hypomania/mania was not assessed at the initial interview in 1979. A formal diagnosis of hypomania based on DSM-IV and strict criteria was applied from the third to the sixth interview. Hypomania was defined in three different ways: (1) DSM-IV criteria, requiring elevated mood plus 3 of 7 criterial symptoms of mania plus a minimum duration of four days; (2) "strict" criteria, requiring 3 of 7 DSM-IV criterial symptoms, but not its time criterion of 4 days, and in addition behavioural consequences (subjectively perceived problems [financial consequences, trouble with the police, etc.] and critical remarks by significant others about a behavioural change); and (3) "broad" criteria, requiring the presence of any hypomanic symptoms without any time restriction and omitting the behavioural consequences. These three sets of criteria for hypomania were used to define three corresponding categories of BP-II: DSM-IV, strict, and broad, all requiring in addition the presence of a MDE. In the last four interviews (1986–1999), in which the number of manic symptoms was assessed, all subjects meeting broad criteria manifested at least 2 of 7 criterial symptoms of mania. Details on the validity of definitions 2 and 3 have been published [5].

Age of onset was evaluated retrospectively after having assessed the whole syndrome and defined as the first manifestation of the syndrome; the latter did not have to qualify for the diagnosis.

Statistical analysis

Prevalence rates and standard errors were computed with adjustment for sample stratification. Lifetime prevalence rates (cumulative lifetime incidence) refer to the total of the six 1-year prevalence rates across the six interviews. Associations between each mood disorder subtype and alcohol abuse and dependence were assessed using logistic regression models, adjusting for stratified sampling and sex. For group comparisons, χ^2 -tests, Fisher's exact tests and Kruskal-Wallis tests were applied. Analyses were conducted using SAS for Windows version 8.01 and Stata 8.0.

Table 1 Design of Zurich cohort study

| Year | Females | Males | Total N | Age (F/M) | Assessment |
|------|---------|-------|---------|-----------|---------------|
| 1978 | 2346 | 2201 | 4547 | 19/20 | Screening |
| 1979 | 299 | 292 | 591 | 20/21 | Interview |
| 1980 | 270 | 234 | 504 | 21/22 | Questionnaire |
| 1981 | 236 | 220 | 456 | 22/23 | Interview |
| 1986 | 232 | 225 | 457 | 27/28 | Interview |
| 1988 | 224 | 200 | 424 | 29/30 | Interview |
| 1993 | 215 | 192 | 407 | 34/35 | Interview |
| 1999 | 205 | 162 | 367 | 40/41 | Interview |

Table 2 Cumulative prevalence rates of alcohol use and BP-II disorders

| | M + F | Males | Females | |
|-----------------------|-------------------------|------------|------------|-----------------------|
| Diagnoses | Prevalence rates % (SE) | | | <i>P</i> (χ^2) |
| Alcohol use disorders | | | | |
| All | 17.9 (2.3) | 28.0 (3.9) | 8.0 (2.2) | 0.000 |
| Dependence | 8.7 (1.6) | 14.5 (3.0) | 3.1 (1.2) | 0.000 |
| Abuse | 9.2 (1.8) | 13.5 (3.0) | 5.0 (1.9) | 0.000 |
| MDE | 22.4 (2.4) | 16.9 (3.1) | 27.7 (3.7) | 0.03 |
| BP-II disorder | | | | |
| DSM-IV | 1.7 (7.6) | 0.3 (0.2) | 2.9 (1.5) | 0.000 |
| Strict | 5.3 (1.3) | 4.1 (1.5) | 6.5 (2.1) | 0.33 |
| Broad | 11.0 (1.8) | 9.2 (2.4) | 12.6 (2.7) | 0.36 |

Results

■ Prevalence of AUD and BP-II

The sex-specific lifetime prevalence rates of AUD and BP-II according to the varying definitions given above are presented in Table 2. Eighteen percent of the sample fulfilled criteria for AUD. The prevalence of BP-II increased monotonically across the 3 increasingly broad sets of criteria, with 1.7% for the DSM-IV criteria, 5.3% for the strict, and 11.0% for the broad criteria. Males had greater rates of both alcohol abuse and dependence than females. By contrast, females had significantly higher rates of DSM-IV MDE and BP-II. The sex difference diminished as the definition of bipolar disorder broadened.

■ Sex, alcohol dependence and alcohol abuse

Table 3 presents the sex-specific associations between AUD and major mood disorders. The association between AUD and mood disorders was consistently stronger in females than in males. This was true for




Fig. 1 Proportion of subjects with alcohol abuse/dependence in mood disorder subgroups based on varying definitions of BP-II disorder. OR = Odds ratio (adjusted for sex and stratified sampling), MDE = Major Depressive Episode, MDD = Major Depressive Disorder, BP = Bipolar Disorder

both alcohol dependence and abuse. For both males and females the association of mood disorders with alcohol dependence was also consistently stronger than with alcohol abuse. For both men and women, the associations between AUD and MDD decreased markedly in step with the broadening of the diagnostic concept of BP-II.

■ AUD according to the varying definitions of BP-II

Figure 1 shows the rates of AUD among subjects with mood disorders, subdivided according to MDE and the varying definitions of bipolar disorder. AUD was much more strongly associated with BP-II than with MDD, irrespective of the definition of bipolar disorder (DSM-IV, strict or broad criteria). Using increasingly broad definitions of bipolar disorder, the association between AUD and MDD decreased from 29.8% for DSM-IV BP-II disorder to 18.8% for the broadest concept of BP-II. With the broadest definition of BP-II, the association between MDD and AUD ceased to be significant

Table 3 Associations (odds ratios = OR) of alcohol abuse, dependence with mood disorders by sex

| Mood disorder | | N | % with alcohol abuse | OR | 95% CI | P | % with alcohol dependence | OR | 95% CI | P |
|---------------|---------------------|-----|----------------------|------------|-----------|-------|---------------------------|------------|-----------|-------|
| Men | MDE ^a | 72 | 15.3 | 1.3 | 0.6–2.6 | 0.581 | 34.7 | 3.8 | 2.0–7.1 | 0.000 |
| | BP-II DSM-IV | 4 | 25.2 | 2.3 | 0.2–22.7 | 0.481 | 75.0 | 5.8 | 1.6–20.2 | 0.006 |
| | Remaining MDE | 68 | 14.7 | 1.2 | 0.5–2.6 | 0.674 | 32.3 | 2.9 | 1.7–4.9 | 0.000 |
| | BP-II strict | 21 | 23.8 | 2.1 | 0.7–6.3 | 0.166 | 38.1 | 5.8 | 2.8–11.8 | 0.000 |
| | Remaining MDE = MDD | 51 | 11.8 | 0.8 | –0.4–2–3 | 0.852 | 33.3 | 2.4 | 1.4–4.2 | 0.002 |
| | BP-II broad | 36 | 19.4 | 1.7 | 0.7–4.1 | 0.281 | 44.4 | 4.8 | 2.7–8.5 | 0.000 |
| | Remaining MDE = MDD | 36 | 11.1 | 0.8 | 0.3–2.6 | 0.786 | 25.0 | 1.9 | 0.95–3–6 | 0.071 |
| Women | MDE | 118 | 6.8 | 2.1 | 0.7–6.3 | 0.174 | 12.7 | 5.2 | 1.8–14.5 | 0.002 |
| | BP-II DSM-IV | 8 | 12.5 | 4.2 | 0.4–39.4 | 0.213 | 12.5 | 5.0 | 0.52–49.0 | 0.164 |
| | Remaining MDE | 110 | 6.4 | 2.0 | 0.6–6.1 | 0.230 | 12.5 | 5.1 | 1.8–14.7 | 0.002 |
| | BP-II strict | 24 | 12.5 | 4.2 | 0.96–17.9 | 0.055 | 29.2 | 14.5 | 4.1–50.6 | 0.000 |
| | Remaining MDE = MDD | 94 | 5.3 | 1.6 | 0.5–5.5 | 0.425 | 8.5 | 3.3 | 1.0–10.3 | 0.043 |
| | BP-II broad | 53 | 13.2 | 4.4 | 1.4–13.8 | 0.010 | 18.9 | 8.2 | 2.7–25.2 | 0.000 |
| | Remaining MDE = MDD | 65 | 1.5 | 0.5 | 0.05–3.8 | 0.471 | 7.7 | 2.9 | 0.8–10.5 | 0.098 |

^aMDE = Major Depression Episode




Fig. 2 Cumulative age of onset of first depressive/manic episode and heavy drinking, for three increasingly broad definitions of BP-II disorder. For each definition, depressive/manic manifestations tend to precede the occurrence of heavy drinking

(OR = 1.0, 95% CI = 0.5, 1.7). Despite the broadening of the concept of bipolarity the association between AUD and BP-II remained absolutely stable (OR = 5.4).

■ Temporal sequence of BP-II and AUD

We compared the ages of the onset of the first depressive/manic episode and the start of heavy drinking in subjects who developed both BP-II and AUD at any later time. This intra-individual comparison was made for the three different definitions of BP-II. Using the broad definition of BP-II, 23 of 34 cases (68%) manifested an affective episode before the onset of heavy drinking; the mean age of onset was 12.9 years (SD 5.03) for bipolarity and 18.2 years (SD 4.19) for heavy drinking. Using the strict definition of BP-II, the first affective episode preceded the onset of heavy drinking in 13 of 19 cases (68%); the mean age of onset was 12.3 years (SD 5.74) for bipolarity and 17.8 years (SD 4.28) for heavy drinking. Finally, all 6 cases with the DSM definition of BP-II (who also had AUD) manifested an affective episode earlier than heavy drinking; the mean age of onset for bipolarity was 9.5 years (SD 6.09), and that for heavy drinking 18.8 years (SD 4.58).

Figure 2 shows the cumulative distribution of ages of onset of the first affective episode and heavy drinking in the three increasingly broad groups of BP-II. In each group, the curve for BP-II is shifted towards earlier ages compared to the curves showing the onset of heavy drinking.

The number of subjects with AUD increased from 14 (at the age of 20–22) to 51 at the age of 40. At the same time, the number of BP-II cases decreased from 47 (at 20–22) to 31 at the age of 40. This finding further supports the view that bipolar symptoms precede and may contribute to the development of AUD.

Discussion

Applying a spectrum concept of BP-II disorder, which resulted in an increasing proportion of MDD subjects being classified as BP-II, we found that the association between MDD and AUD progressively diminished and ultimately disappeared with the broadest definition of BP-II. Whereas 93.5% of AUD cases had MDD according to DSM-IV criteria, the figure was only 53% of the AUD cases had MDD with our broadest definition of bipolar disorder. Our findings also confirm that the association between alcoholism and mood disorders was far greater for alcohol dependence than for abuse, particularly in men. This supports prior family study research suggesting that women have a lower threshold for expression of alcoholism than men [24].

Another key finding of our study relates to the time sequence of the onset of symptoms of AUD and BP-II. Affective episodes preceded heavy drinking in most cases. In the majority of cases the onset of BP occurred before age 20, whereas at this age AUD was a minor problem but grew quickly in the third decade of life. In the prospective period from age 20/21 to 40/41 changes from BP-II to AUD were several times more frequent than the reverse (data not shown).

These findings could have major implications for risk factor research and the prevention of AUD. If our findings are replicable and, in particular, if BP-II disorder tends to manifest first in the comorbid sequence, this could have a substantial impact on the prevention of AUD, especially in women. Application of broad rather than strict diagnostic criteria would allow earlier and more accurate identification of individuals at risk. The administration of instruments for the self-assessment of hypomania/mania in cases of depression [4, 18] could greatly help to identify bipolar subjects, to initiate early prophylactic treatment and to reduce their risk of developing secondary alcohol and drug abuse/dependence.

In terms of clinical implications, our findings have relevance for both diagnosis and treatment. The finding that AUD are predominantly associated with bipolar disorder suggests the need to screen for bipolarity in all subjects with alcohol problems. Such screening could also apply in the evaluation of dual diagnoses in people with schizophrenia or schizo-affective with substance use disorders [9, 25, 29]. Increased vigilance in the detection of bipolarity among those who have already developed dependence, could be used to select treatment modalities that address both the mood and substance problems. Indeed, there is emerging evidence for the success of this approach in a growing number of studies [8, 13]. Future studies will be required to examine the extent to which data from the present study have relevance for intervention in the transition to early and mid adulthood.

Finally, these findings have important implications for etiologic studies; particularly those which seek to identify genetic factors underlying mood disorders or alcoholism. Although evidence regarding the extent to which comorbidity between alcoholism and bipolar disorder is attributable to share common underlying genetic risk factors is inconclusive [6, 22, 24], none of the prior family and genetic studies employed the spectrum concept of bipolar disorders [14]. Therefore, results of earlier studies may have underestimated the extent to which bipolar disorder and alcoholism have familial links.

The findings of this study should be considered in the context of the following limitations: (a) subjects were aged 20/21 when the study started; the data on age of onset are therefore partially retrospective; (b) differences in diagnostic information available across six interview waves (DSM-III and DSM-III-R major depression); (c) the 20-year attrition rate of 38%; (d) the relatively small sample size of this cohort; which *inter alia* did not allow assessment of rare disorders.

■ **Acknowledgement** This work was supported by Grant 3200-050881.97/1 of the Swiss National Science Foundation

References

1. Akiskal H, Bourgeois ML, Angst J, Post R, Möller HJ, Hirschfeld R (2000) Re-evaluating the prevalence of and diagnostic composition within the broad clinical spectrum of bipolar disorders. *J Affect Disord* 59:S5–S30
2. Allgood-Merten B, Lewinsohn PM, Hops H (1990) Sex differences and adolescent depression. *J Abnorm Psychol* 99:55–63
3. Angst J (1992) L'hypomanie. A propos d'une cohorte de jeunes. *Encephale* 18:23–29
4. Angst J, Adolfsson R, Benazzi F, Gamma A, Hantouche E, Meyer TD, Skeppar P, Vieta E, Scott J (2005) The HCL-32: towards a self-assessment tool for hypomanic symptoms in outpatients. *J Affect Disord* 88:217–233
5. Angst J, Gamma A, Benazzi F, Ajdacic V, Eich D, Rössler W (2003) Toward a re-definition of subthreshold bipolarity: epidemiology and proposed criteria for bipolar-II, minor bipolar disorders and hypomania. *J Affect Disord* 73:133–146
6. Bierut LJ, Saccone NL, Price JP, Goate A, Foroud T, Edenberg H, Almasy L, Conneally PM, Crowe R, Hesselbrock V, Li TK, Nurnberger J, Porjesz B, Schuckit MA, Tischfield J, Begleiter H, Reich T (2002) Defining alcohol-related phenotypes in humans. *Alcohol Res Health* 26:208–213
7. Brady KT, Lydiard RB (1992) Bipolar affective disorder and substance abuse. *J Clin Psychopharmacol* 12:17S–22S
8. Brown ES (2005) Bipolar disorder and substance abuse. *Psychiatr Clin N Am* 28:415–425
9. Caspari D (1999) Cannabis and schizophrenia: results of a follow-up study. *Eur Arch Psychiatr Neurol Sci* 249:45–49
10. Chengappa KNR, Levine J, Gershon S, Kupfer DJ (2000) Lifetime prevalence of substance or alcohol abuse and dependence among subjects with bipolar I and II disorders in a voluntary registry. *Bipolar Disord* 2:191–195
11. Derogatis LR, Cleary PA (1977) Factorial invariance across gender for the primary symptom dimensions of the SCL-90. *Br J Soc Clin Psychol* 16:347–356
12. Fogarty F, Russell JM, Newman SC, Bland RC (1994) Mania. *Acta Psychiatr Scand* 376:16–23
13. Geller B, Cooper TB, Sun K, Zimmerman B, Frazier J, Williams M, Heath J (1998) Double-blind and placebo-controlled study of lithium for adolescent bipolar disorders with secondary substance dependency. *J Am Acad Child Adolesc Psychiatry* 37:171–178
14. Ghaemi N, Sachs GS, Goodwin FK (2000) What is to be done? Controversies in the diagnosis and treatment of manic-depressive illness. *World J Psychiatry* 1:65–74
15. Grant BF, Dawson DA, Stinson FS, Chou PS, Kay W, Pickering R (2003) The Alcohol Use Disorder and Associated Disabilities Interview Schedule-IV (AUDADIS-IV): reliability of alcohol consumption, tobacco use, family history of depression and psychiatric diagnostic modules in a general population sample. *Drug Alcohol Depend* 71:7–16
16. Grant BF, Stinson FS, Dawson DA, Chou SP, Dufour MC, Compton W, Pickering RP, Kaplan K (2004) Prevalence and co-occurrence of substance use disorders and independent mood and anxiety disorders. Results from the National Epidemiologic Survey on alcohol and related conditions. *Arch Gen Psychiatry* 61:807–816
17. Hantouche EG, Akiskal HS, Lancrenon S, Allilaire J-F, Sechter D, Azorin J-M, Bourgeois M, Fraud J-P, Châtenet-Duchêne L (1998) Systematic clinical methodology for validating bipolar - II disorder: data in mid-stream from a French national multi-site study (EPIDEP). *J Affect Disord* 50:163–173
18. Hirschfeld RMA, Williams JBW, Spitzer RL, Calabrese JR, Flynn L, Keck P, Lewis L, McElroy S, Post RM, Rappaport DJ (2000) Development and validation of a screening instrument for bipolar spectrum disorder: the mood disorder questionnaire. *Am J Psychiatry* 157:1873–1875
19. Kessler RC, Crum RM, Warner LA, Nelson CB, Schulenberg J, Anthony JC (1997) Lifetime co-occurrence of DSM-III-R alcohol abuse and dependence with other psychiatric disorders in the National Comorbidity Survey. *Arch Gen Psychiatry* 54:313–321
20. Lewinsohn PM, Klein DN, Seeley JR (2000) Bipolar disorder during adolescence and young adulthood in a community sample. *Bipolar Disord* 2:281–293
21. Lewinsohn PM, Klein DN, Seeley JR (1995) Bipolar disorders in a community sample of older adolescents: prevalence, phenomenology, comorbidity, and course. *J Am Acad Child Adolesc Psychiatry* 34:454–463
22. Maier W, Merikangas KR (1992) Contranmission and comorbidity of affective disorders, anxiety disorders and alcoholism in families. In: Mendlewicz J, Hippus H (eds) *Genetic research in psychiatry*. Springer, New York, pp 210–224
23. Merikangas KR, Angst J, Eaton W, Canino G, Rubio-Stipec M, Wacker H, Wittchen HU, Andrade L, Essau C, Whitaker A, Kraemer H, Robins LN, Kupfer DJ (1996) Comorbidity and boundaries of affective disorders with anxiety disorders and substance misuse: results of an international task force. *Br J Psychiatry* 168:58–67
24. Merikangas KR, Stevens DE, Fenton B, Stolar M, O'Malley S, Woods SW, Risch N (1998) Co-morbidity and familial aggregation of alcoholism and anxiety disorders. *Psychol Med* 28:773–788
25. Modestin J, Nussbaumer C, Angst K, Scheidegger P, Hell D (1997) Use of potentially abusive psychotropic substances in psychiatric inpatients. *Eur Arch Psychiatry Clin Neurosci* 247:146–153
26. Preuss UW, Schickit MA, Smith TL, Danko GP, Buckman K, Bierut L, Bucholz KK, Hesselbrock MN, Hesselbrock VM, Reich T (2002) Comparison of 3190 alcohol-dependent individuals with and without suicide attempts. *Alcohol Clin Exp Res* 26:471–477
27. Preuss UW, Schuckit MA, Smith TL, Danko GP, Dasher AC, Hesselbrock MN, Hesselbrock VM, Nurnberger JI (2002) A comparison of alcohol-induced and independent depression in alcoholics with histories of suicide attempts. *J Stud Alcohol* 63:498–502
28. Regier DA, Farmer ME, Rae DS, Locke BZ, Keith SJ, Judd LL, Goodwin FK (1990) Comorbidity of mental disorders with alcohol and other drug abuse. Results from the Epidemiologic Catchment Area (ECA) study. *JAMA* 264:2511–2518

29. Soyka M, Albus M, Immler B, Kathmann N, Hippus H (2001) Psychopathology in dual diagnosis and non-addicted schizophrenics—are there differences? *Eur Arch Psychiatr Neurol Sci* 251:232–238
30. Winokur G, Turvey C, Akiskal H, Coryell W, Solomon D, Leon A, Mueller T, Endicott J, Maser J, Keller M (1998) Alcoholism and drug abuse in three groups—bipolar I, unipolars and their acquaintances. *J Affect Disord* 50:81–89